

A QUANTUM ANALOGUE OF THE BERNSTEIN FUNCTOR

S. Sinel'shchikov[†]

A. Stolin[‡]

L. Vaksman[†]

[†]*Mathematics Division, Institute for Low Temperature Physics & Engineering
47 Lenin Ave, 61103 Kharkov, Ukraine*

[‡]*Chalmers Tekniska Högskola, Matematik, 412 96, Göteborg, Sweden*

Abstract

We consider Knapp-Vogan Hecke algebras in the quantum group setting. This allows us to produce a quantum analogue of the Bernstein functor as a first step towards the cohomological induction for quantum groups.

The cohomological induction is one of the most important tools for producing unitarizable Harish-Chandra modules [7]. The present paper demonstrates that this method can be applied to modules over quantum universal enveloping algebras [6] as well. Our way to this circle of problems was as follows.

The papers [11, 9] introduce quantum analogues of the bounded symmetric domains and the associated Harish-Chandra modules. One special case considered in [10] leads to two geometric realizations for the ladder representation of the quantum universal enveloping algebra $U_q \mathfrak{su}_{2,2}$ and to the quantum Penrose transform [1]. One of the main tools used in [10] was studying q -analogues of the Čech cohomology. Unfortunately this method can be hardly generalized because it relates to non-trivial problems in non-commutative algebraic geometry. We hope to overcome these obstacles via replacing the q -analogues of the Čech cohomology by q -analogues of the Dolbeault cohomology. It is well-known that in the classical situation the Dolbeault cohomology can be constructed algebraically using the so-called cohomological induction (see [7]).

It is convenient to work with the Bernstein functor (the projective Zuckerman functor) for producing unitarizable Harish-Chandra modules. The definition of the Bernstein functor uses essentially the algebra of distributions on a real reductive group G with support in a maximal compact subgroup K . The principal obstacle to cope with in this work is to construct a q -analogue for this algebra.

In the present paper we construct a q -analogue of the algebra of distributions as above in certain cases. This allows us to construct a q -analogue of the Bernstein functor.

The authors are grateful to V. Kruglov for assistance in preparation of a preliminary version of this work and for helpful discussions.

1 The category $C(\mathfrak{l}, \mathfrak{l})_q$ and the algebra $R(\mathfrak{l})_q$

We will use $\mathbb{C}$ as the ground field.

Let $\mathbf{a} = (a_{ij})_{i,j=1,2,\dots,l}$, be the Cartan matrix for a simple complex Lie algebra of rank l . Its universal enveloping algebra is a unital algebra U determined by the generators $\{H_i, E_i, F_i\}_{i,j=1,2,\dots,l}$ and the relations [6, p. 51]:

$$\begin{aligned} H_i H_j - H_j H_i &= 0, & E_i F_j - F_j E_i &= \delta_{ij} H_i, \\ H_i E_j - E_j H_i &= a_{ij} E_j, & H_i F_j - F_j H_i &= -a_{ij} F_j, & i, j &= 1, 2, \dots, l, \end{aligned}$$

and

$$\begin{aligned} \sum_{m=0}^{1-a_{ij}} (-1)^m \binom{1-a_{ij}}{m} E_i^{1-a_{ij}-m} \cdot E_j \cdot E_i^m &= 0, \\ \sum_{m=0}^{1-a_{ij}} (-1)^m \binom{1-a_{ij}}{m} F_i^{1-a_{ij}-m} \cdot F_j \cdot F_i^m &= 0, \end{aligned}$$

for all $i \neq j$.

The Cartan subalgebra spanned by $H_1, H_2, \dots, H_l$ is denoted by $\mathfrak{h}$, and $\alpha_1, \alpha_2, \dots, \alpha_l$,

$$\alpha_j(H_i) = a_{ij}, \quad i, j = 1, 2, \dots, l,$$

are simple roots.

There exists a unique collection of coprime positive integers $d_1, d_2, \dots, d_l$ such that

$$d_i a_{ij} = d_j a_{ji}, \quad i, j = 1, 2, \dots, l.$$

The bilinear form in $\mathfrak{h}^*$ given by

$$(\alpha_i, \alpha_j) = d_i a_{ij}, \quad i, j = 1, 2, \dots, l,$$

is positive definite.

Recall the definition of the quantum universal enveloping algebra U_q introduced by V. Drinfeld and M. Jimbo. We assume that $q \in (0, 1)$.¹

The unital algebra U_q is determined by its generators $K_i, K_i^{-1}, E_i, F_i, i = 1, 2, \dots, l$, and the relations

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1, \\ K_i E_j &= q_i^{a_{ij}} E_j K_i, & K_i F_j &= q_i^{-a_{ij}} F_j K_i, & E_i F_j - F_j E_i &= \delta_{ij} (K_i - K_i^{-1}) / (q_i - q_i^{-1}), \\ \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1-a_{ij} \\ s \end{bmatrix}_{q_i} E_i^{1-a_{ij}-s} E_j E_i^s &= 0, \\ \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1-a_{ij} \\ s \end{bmatrix}_{q_i} F_i^{1-a_{ij}-s} F_j F_i^s &= 0, \end{aligned}$$

with

$$q_i = q^{d_i}, \quad 1 \leq i \leq l,$$

$$\begin{bmatrix} m \\ n \end{bmatrix}_q = \frac{[m]_q!}{[n]_q! [m-n]_q!}, \quad [n]_q! = [n]_q \dots [2]_q [1]_q, \quad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

¹The purpose of so strong assumptions on q will become evident while investigating the unitarizability of the Harish-Chandra modules in question over U_q .

The Hopf algebra structure on U_q is given by the following formulas:

$$\begin{aligned}\Delta(E_i) &= E_i \otimes 1 + K_i \otimes E_i, & \Delta(F_i) &= F_i \otimes K_i^{-1} + 1 \otimes F_i, & \Delta(K_i) &= K_i \otimes K_i, \\ S(E_i) &= -K_i^{-1} E_i, & S(F_i) &= -F_i K_i, & S(K_i) &= K_i^{-1}, \\ \varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= 1.\end{aligned}$$

Any subset $\mathbb{L} \subset \{1, 2, \dots, l\}$ determines a Hopf subalgebra $U_q \mathfrak{l} \subset U_q$ generated by

$$K_i^{\pm 1}, \quad i = 1, 2, \dots, l; \quad E_j, F_j, \quad j \in \mathbb{L}.$$

We use the following notation:

$$P = \mathbb{Z}^l, \quad P_+ = \mathbb{Z}_+^l, \quad P_+^{\mathbb{L}} = \{\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P \mid \lambda_i \geq 0 \text{ for } i \in \mathbb{L}\}.$$

Let V be a $U_q \mathfrak{l}$ -module and

$$V_\mu = \{v \in V \mid K_i v = q_i^{\mu_i} v, \quad i = 1, 2, \dots, l\}$$

the weight space, where $\mu = (\mu_1, \mu_2, \dots, \mu_l) \in P$.

Definition 1.1 We say V is a **weight** $U_q \mathfrak{l}$ -module if $V = \bigoplus_{\mu \in P} V_\mu$.

Example 1.2 Consider the Verma module $U_q \mathfrak{l}$ -module $M(\mathfrak{l}, \lambda)$ with the highest weight $\lambda \in P_+^{\mathbb{L}}$. This is a module determined by its generator $v(\mathfrak{l}, \lambda)$ and the relations

$$E_j v(\mathfrak{l}, \lambda) = 0, \quad j \in \mathbb{L}; \quad K_i^{\pm 1} v(\mathfrak{l}, \lambda) = q_i^{\pm \lambda_i} v(\mathfrak{l}, \lambda), \quad i = 1, 2, \dots, l.$$

It is easy to prove that its submodule $K(\mathfrak{l}, \lambda)$ generated by $F_j^{\lambda_j+1} v(\mathfrak{l}, \lambda)$, $j \in \mathbb{L}$, is the unique submodule of finite codimension. In particular, the factor-module $L(\mathfrak{l}, \lambda) = M(\mathfrak{l}, \lambda)/K(\mathfrak{l}, \lambda)$ is simple (see [6]).

The following claims are well-known (see for instance [6]). The dimensions of the weight subspaces of the $U_q \mathfrak{l}$ -modules $L(\mathfrak{l}, \lambda)$ are the same as in the classical case $q = 1$. The weight finite dimensional $U_q \mathfrak{l}$ -modules are completely reducible. Irreducible weight finite dimensional $U_q \mathfrak{l}$ -modules are isomorphic to $L(\mathfrak{l}, \lambda)$ with $\lambda \in P_+^{\mathbb{L}}$. The center of $U_q \mathfrak{l}$ (denoted by $Z(U_q \mathfrak{l})$) admits an easy description via the Harish-Chandra isomorphism [6, p. 109] and separates the simple weight finite dimensional $U_q \mathfrak{l}$ -modules [6, p. 125]. In other words, for any $\lambda_1, \lambda_2 \in P_+^{\mathbb{L}}$ there exists $z \in Z(U_q \mathfrak{l})$ such that $z|_{L(\mathfrak{l}, \lambda_1)} \neq z|_{L(\mathfrak{l}, \lambda_2)}$.

A module V over $U_q \mathfrak{l}$ is called locally finite dimensional if $\dim(U_q \mathfrak{l} v) < \infty$ for all $v \in V$. Introduce the notation $C(\mathfrak{l}, \mathfrak{l})_q$ for the full subcategory of weight locally finite dimensional $U_q \mathfrak{l}$ -modules.

It is easy to prove that any $U_q \mathfrak{l}$ -module $V \in C(\mathfrak{l}, \mathfrak{l})_q$ is uniquely decomposable as a direct sum $V = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} V_\lambda$ of its submodules. Each V_λ here is a multiple of $L(\mathfrak{l}, \lambda)$, $\lambda \in P_+^{\mathbb{L}}$.

The sum is direct since $Z(U_q \mathfrak{l})$ separates simple finite dimensional weight $U_q \mathfrak{l}$ -modules.

Let $\text{End } V$ be the algebra of all endomorphisms of V considered as a vector space. Now consider the $U_q \mathfrak{l}$ -module $L^{\text{univ}} = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} L(\mathfrak{l}, \lambda)$ and the projections P_λ in L^{univ} onto

the isotypic component $L(\mathfrak{l}, \lambda)$ parallel to the sum of all other isotypic components. The

algebra $\text{End } L^{\text{univ}}$ is a $U_q\mathfrak{l}$ -module algebra.² The natural homomorphism $U_q\mathfrak{l} \rightarrow \text{End } L^{\text{univ}}$ is injective [6, p. 76 – 77]; this allows one to identify $U_q\mathfrak{l}$ with its image in $\text{End } L^{\text{univ}}$. We are interested in the following $U_q\mathfrak{l}$ -module subalgebras of $\text{End } L^{\text{univ}}$:

$$R(\mathfrak{l})_q \stackrel{\text{def}}{=} \bigoplus_{\lambda \in P_+^{\mathbb{L}}} \text{End } L(\mathfrak{l}, \lambda), \quad F(\mathfrak{l})_q \stackrel{\text{def}}{=} R(\mathfrak{l})_q \oplus U_q\mathfrak{l}.$$

Obviously, $R(\mathfrak{l})_q$ is a two-sided ideal of $F(\mathfrak{l})_q$ generated by the projections P_λ .

$R(\mathfrak{l})_q$ is not a unital algebra, but it admits a distinguished approximate identity. Specifically, any finite subset $\Lambda \subset P_+^{\mathbb{L}}$ determines an idempotent $\chi_\Lambda = \sum_{\lambda \in \Lambda} P_\lambda$ in $R(\mathfrak{l})_q$. Obviously, $\chi_{\Lambda_1}\chi_{\Lambda_2} = \chi_{\Lambda_2}\chi_{\Lambda_1} = \chi_{\Lambda_1 \cap \Lambda_2}$, and for any $r \in R(\mathfrak{l})_q$ there exists a finite subset $\Lambda \subset P_+^{\mathbb{L}}$ such that $\chi_\Lambda r = r\chi_\Lambda = r$. With some abuse of terminology, one can say that in $F(\mathfrak{l})_q$ we have

$$\lim_{\Lambda \uparrow P_+^{\mathbb{L}}} \chi_\Lambda = 1.$$

A module V over $R(\mathfrak{l})_q$ is called approximately unital if for any $v \in V$ there exists a finite subset $\Lambda \subset P_+^{\mathbb{L}}$ such that $\chi_\Lambda v = v$.

We are going to demonstrate that the full subcategory of approximately unital $R(\mathfrak{l})_q$ -modules is canonically isomorphic to $C(\mathfrak{l}, \mathfrak{l})_q$.

Proposition 1.3 (*). *For any approximately unital $R(\mathfrak{l})_q$ -module V and any $\xi \in F(\mathfrak{l})_q$, $v \in V$ there exists a limit*

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_+^{\mathbb{L}}} (\xi \chi_\Lambda) v \tag{1.1}$$

in the sense that $(\xi \chi_\Lambda) v$ does not depend on Λ for large Λ .

(**). *The relation (1.1) equips V with a structure of $F(\mathfrak{l})_q$ -module and defines a functor from the category of approximately unital $R(\mathfrak{l})_q$ -modules into the category $C(\mathfrak{l}, \mathfrak{l})_q$.*

(***). *This functor is an isomorphism of categories.*

Proof. Any approximately unital $R(\mathfrak{l})_q$ -module V admits an embedding into an approximately unital $R(\mathfrak{l})_q$ -module that is a multiple of L^{univ} . This observation implies the first two statements, since they can be easily verified for L^{univ} .

Any $U_q\mathfrak{l}$ -module that is a multiple of L^{univ} , is a $F(\mathfrak{l})_q$ -module. This allows one to construct an inverse functor by embedding any $U_q\mathfrak{l}$ -module $V \in C(\mathfrak{l}, \mathfrak{l})_q$ into a $U_q\mathfrak{l}$ -module that is a multiple of L^{univ} . This implies (***). $\square$

Note that the action of $P_\lambda \in R(\mathfrak{l})_q$ on vectors $v \in V \in C(\mathfrak{l}, \mathfrak{l})_q$ can be described without referring to an embedding into a $U_q\mathfrak{l}$ -module multiple to L^{univ} . More exactly, for any $\lambda \in P_+^{\mathbb{L}}$ there exists a sequence $z_1(\lambda), z_2(\lambda), \dots$ of elements of the center of $U_q\mathfrak{l}$ such that $P_\lambda v = \lim_{n \rightarrow \infty} z_n(\lambda) v$ for all $v \in V \in C(\mathfrak{l}, \mathfrak{l})_q$. To see this, note that for any finite subset $\Lambda \subset P_+^{\mathbb{L}}$ there exists such $z \in Z(U_q\mathfrak{l})$ that $z|_{L(\mathfrak{l}, \mu)} = 0$ for $\lambda \in \Lambda \setminus \{\lambda\}$ and $z|_{L(\mathfrak{l}, \lambda)} = 1$. It remains to choose a sequence $\Lambda_n \uparrow P_+^{\mathbb{L}}$.

²If $\xi \in U_q\mathfrak{l}$ and $\Delta \xi = \sum_i \xi'_i \otimes \xi''_i$, then $(\xi A)v = \sum_i \xi'_i A S(\xi''_i) v$, $v \in L^{\text{univ}}$, $A \in \text{End } L^{\text{univ}}$, with Δ and S being the comultiplication and the antipode of $U_q\mathfrak{l}$.

³It is easy to prove that $R(\mathfrak{l})_q \cap U_q\mathfrak{l} = 0$. In fact, $U_q\mathfrak{l}$ has no zero divisors [3], while the Harish-Chandra isomorphism allows one to produce for any $a \in R(\mathfrak{l})_q$ a non-zero element $z \in Z(U_q\mathfrak{l})$ such that $za = 0$.

2 The category $C(\mathfrak{g}, \mathfrak{l})_q$ and the algebra $R(\mathfrak{g}, \mathfrak{l})_q$

Let $\mathbb{G} \supset \mathbb{L}$ be a pair of subsets of $\{1, 2, \dots, l\}$ and let $U_q \mathfrak{g} \supset U_q \mathfrak{l}$ be the associated pair of Hopf subalgebras of U_q .

Consider the category of all $U_q \mathfrak{g}$ -modules and its full subcategory $C(\mathfrak{g}, \mathfrak{l})_q$ that consists of weight $U_q \mathfrak{l}$ -locally finite dimensional $U_q \mathfrak{g}$ -modules.

Let us turn to a description of the algebra $F(\mathfrak{g}, \mathfrak{l})_q$. This algebra is an important tool in studying $U_q \mathfrak{g}$ -modules from $C(\mathfrak{g}, \mathfrak{l})_q$. First, we consider a simple special case, namely $\mathbb{G} = \mathbb{L}$. In this situation we define $F(\mathfrak{g}, \mathfrak{l})_q$ as $F(\mathfrak{l})_q$. We intend to construct $F(\mathfrak{g}, \mathfrak{l})_q$ along with embeddings $U_q \mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, which possess the following properties.

(F1) The diagram

$$\begin{array}{ccc} U_q \mathfrak{l} & \xrightarrow{\quad} & U_q \mathfrak{g} \\ \downarrow & & \downarrow \\ F(\mathfrak{l})_q & \xrightarrow{\quad} & F(\mathfrak{g}, \mathfrak{l})_q \end{array}$$

commutes⁴, and the subalgebras $U_q \mathfrak{g}$ and $F(\mathfrak{l})_q$ generate the algebra $F(\mathfrak{g}, \mathfrak{l})_q$.

(F2) The approximate identity $\{\chi_\Lambda\}$ of $R(\mathfrak{l})_q$ is also an approximate identity of the two-sided ideal $R(\mathfrak{g}, \mathfrak{l})_q$ of $F(\mathfrak{g}, \mathfrak{l})_q$, generated by $R(\mathfrak{l})_q$.

(F3) For any approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -module V and any $\xi \in F(\mathfrak{g}, \mathfrak{l})_q$, $v \in V$ there exists a limit

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_+^L} (\xi \chi_\Lambda) v, \quad (2.1)$$

and the equation (2.1) equips V with a structure of $F(\mathfrak{g}, \mathfrak{l})_q$ -module.

The intersection of the kernels for all representations of $F(\mathfrak{g}, \mathfrak{l})_q$ obtained in this way is 0.

(F4) The functor from the category of approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -modules into $C(\mathfrak{g}, \mathfrak{l})_q$ given by (2.1) is an isomorphism of categories.

We are going to show that the algebra $F(\mathfrak{g}, \mathfrak{l})_q$ and the embeddings $U_q \mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ are essentially uniquely determined by (F1) – (F4), i.e. there exists a unique isomorphism of algebras that respects the embeddings.

Assign to any $U_q \mathfrak{l}$ -module V a $U_q \mathfrak{g}$ -module $P(V) = U_q \mathfrak{g} \otimes_{U_q \mathfrak{l}} V$. The following result is well-known in the classical case $q = 1$ (see [7]). It is proved in the Appendix (see Corollary A.3).

Lemma 2.1 1. If $V \in C(\mathfrak{l}, \mathfrak{l})_q$ then $P(V)$ is a projective object in $C(\mathfrak{g}, \mathfrak{l})_q$.

2. For any $V \in C(\mathfrak{g}, \mathfrak{l})_q$, the map

$$P(V) \mapsto V, \quad \xi \otimes v \mapsto \xi v, \quad \xi \in U_q \mathfrak{g}, \quad v \in V,$$

is a surjective morphism of $U_q \mathfrak{g}$ -modules.

⁴In what follows $U_q \mathfrak{g}$ and $F(\mathfrak{l})_q$ are identified to their images under the embedding into $F(\mathfrak{g}, \mathfrak{l})_q$.

The projective objects $P(V)$ of the category $C(\mathfrak{g}, \mathfrak{l})_q$ are called standard projective objects.

Introduce the $U_q\mathfrak{g}$ -module $V^{\text{univ}} = P(L^{\text{univ}})$.

Corollary 2.2 *In the category of $U_q\mathfrak{l}$ -modules one has*

$$V^{\text{univ}} = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} V_{\lambda}^{\text{univ}},$$

where the $U_q\mathfrak{l}$ -modules $V_{\lambda}^{\text{univ}}$ are multiples of $L(\mathfrak{l}, \lambda)$.

Corollary 2.3 *For any module $M \in C(\mathfrak{g}, \mathfrak{l})_q$ there exists a surjective morphism $\tilde{V}^{\text{univ}} \rightarrow M$ in the category $C(\mathfrak{g}, \mathfrak{l})_q$, with $\tilde{V}^{\text{univ}}$ being a multiple of the module V^{univ} .*

Let us introduce the notation $\mathcal{P}_{\lambda}$ for the projection in V^{univ} onto the $U_q\mathfrak{l}$ -isotypic component $V_{\lambda}^{\text{univ}}$ parallel to the sum of all other $U_q\mathfrak{l}$ -isotypic components.

The category $C(\mathfrak{g}, \mathfrak{l})_q$ and the category of approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -modules are closed under the operations of direct sums, passage to submodules and factor-modules. This allows one, using Lemma 2.1 and Corollary 2.3, to prove the following statement, which implies uniqueness of $F(\mathfrak{g}, \mathfrak{l})_q$ and the embeddings $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$.

Proposition 2.4 *Consider the algebra $F(\mathfrak{g}, \mathfrak{l})_q$ and the embeddings $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ with the properties (F1) – (F4). Then the following statements hold.*

1. *The representation of $U_q\mathfrak{g}$ in V^{univ} determines an approximately unital representation of $R(\mathfrak{g}, \mathfrak{l})_q$ and a faithful representation of $F(\mathfrak{g}, \mathfrak{l})_q$ in V^{univ} .*
2. *The image of $F(\mathfrak{g}, \mathfrak{l})_q$ under the embedding into $\text{End } V^{\text{univ}}$ is the subalgebra of $\text{End } V^{\text{univ}}$ generated by elements of $U_q\mathfrak{g} \subset \text{End } V^{\text{univ}}$ and projections $\mathcal{P}_{\lambda}$, $\lambda \in P_+^{\mathbb{L}}$.*

Now we return to the construction of the algebra $F(\mathfrak{g}, \mathfrak{l})_q$. In what follows we will identify the algebras $U_q\mathfrak{g}$, $F(\mathfrak{l})_q$ with their images under the embeddings into $\text{End } V^{\text{univ}}$.⁵ In particular,

$$P_{\lambda} \mapsto \mathcal{P}_{\lambda}, \quad \lambda \in P_+^{\mathbb{L}}.$$

The following auxiliary statement is proved in the Appendix.

Lemma 2.5 *Given any $\xi \in U_q\mathfrak{g}$, $\lambda \in P_+^{\mathbb{L}}$, there exists a finite subset $\Lambda \subset P_+^{\mathbb{L}}$ such that*

$$\xi \mathcal{P}_{\lambda} = \chi_{\Lambda} \xi \mathcal{P}_{\lambda}, \quad \mathcal{P}_{\lambda} \xi = \mathcal{P}_{\lambda} \xi \chi_{\Lambda}.$$

Consider the smallest subalgebra $F(\mathfrak{g}, \mathfrak{l})_q \subset \text{End } V^{\text{univ}}$ that contains all the elements of $U_q\mathfrak{g}$ and all the projections $\mathcal{P}_{\lambda}$, $\lambda \in P_+^{\mathbb{L}}$.

Theorem 2.6 *The algebra $F(\mathfrak{g}, \mathfrak{l})_q$ along with the embeddings $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$, satisfy (F1) – (F4).*

⁵The representation of $U_q\mathfrak{g}$ in V^{univ} is faithful since for any weight finite dimensional $U_q\mathfrak{g}$ -module W there exists a surjective morphism of $U_q\mathfrak{g}$ -modules $V^{\text{univ}} \rightarrow W$ (Corollary 2.3). The representation of $F(\mathfrak{l})_q$ in V^{univ} is faithful since $L^{\text{univ}} \hookrightarrow V^{\text{univ}}$.

Proof. (F1) is obviously satisfied.

Let us prove that $\{\chi_\Lambda\}$ is a left approximate identity in $R(\mathfrak{g}, \mathfrak{l})_q$. Every $a \in R(\mathfrak{g}, \mathfrak{l})_q$ has the form $a = \sum_{i=1}^{N(a)} \xi_i \mathcal{P}_{\lambda_i} \eta_i$, with $\xi_i \in U_q \mathfrak{g}$, $\eta_i \in F(\mathfrak{g}, \mathfrak{l})_q$, $\lambda_i \in P_+^\mathbb{L}$. It follows from Lemma 2.5 that $\chi_\Lambda \xi_i \mathcal{P}_{\lambda_i} = \xi_i \mathcal{P}_{\lambda_i}$ for some finite subset $\Lambda \subset P_+^\mathbb{L}$ and all $i = 1, 2, \dots, N(a)$. Hence $\chi_\Lambda a = a$. One can prove in a similar way that $\{\chi_\Lambda\}$ is a right approximate identity in $R(\mathfrak{g}, \mathfrak{l})_q$. Hence (F2) is satisfied.

Consider an approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -module V . Given any vector $v \in V$, there exists such finite subset $\Lambda \subset P_+^\mathbb{L}$ that $\chi_\Lambda v = v$. Thus for all $\Lambda' \supset \Lambda$ we have

$$(\xi \chi_{\Lambda'})v = (\xi \chi_{\Lambda'})\chi_\Lambda v = (\xi \chi_{\Lambda'} \chi_\Lambda)v = (\xi \chi_\Lambda)v,$$

and hence the element

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_+^\mathbb{L}} (\xi \chi_\Lambda)v$$

is well defined. It is easy to prove that $(\xi \eta)v = \xi(\eta v)$ for any $\xi, \eta \in F(\mathfrak{g}, \mathfrak{l})_q$, $v \in V$. In fact, there exist such finite subsets $\Lambda', \Lambda'' \subset P_+^\mathbb{L}$ that

$$\chi_{\Lambda'} v = v, \quad \chi_{\Lambda''} \eta \chi_{\Lambda'} = \eta \chi_{\Lambda'},$$

because $\eta \chi_{\Lambda'} \in R(\mathfrak{g}, \mathfrak{l})_q$ and $\{\chi_\Lambda\}$ is an approximate identity. Hence,

$$\begin{aligned} (\xi \eta)v &= \lim_{\Lambda \uparrow P_+^\mathbb{L}} (\xi \eta \chi_\Lambda)v = \xi(\eta \chi_{\Lambda'})v = (\xi \chi_{\Lambda''})(\eta \chi_{\Lambda'})v = \lim_{\Lambda \uparrow P_+^\mathbb{L}} (\xi \chi_\Lambda)(\eta \chi_{\Lambda'})v = \\ &= \xi(\eta \chi_{\Lambda'} v) = \xi(\eta v). \end{aligned}$$

It follows that V is an $F(\mathfrak{g}, \mathfrak{l})_q$ -module, and the first of the conditions (F3) is satisfied. The second condition, which requires the existence of a faithful representation as above, obviously holds.

The arguments below show that the $F(\mathfrak{g}, \mathfrak{l})_q$ -modules in question are locally $U_q \mathfrak{l}$ -finite dimensional and weight:

$$\dim(U_q \mathfrak{l} v) = \dim U_q \mathfrak{l}(\chi_{\Lambda'} v) = \dim(U_q \mathfrak{l} \chi_{\Lambda'} v) \leq \dim R(\mathfrak{l})_q v < \infty.$$

Thus we have constructed a functor from the category of approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -modules into $C(\mathfrak{g}, \mathfrak{l})_q$. What remains is to construct an inverse functor.

Suppose $V \in C(\mathfrak{g}, \mathfrak{l})_q$ and let $V = \bigoplus_{\mu \in P_+^\mathbb{L}} V_\mu$ be a decomposition of V into a sum of

$U_q \mathfrak{l}$ -isotypic components. Let π be the representation of $U_q \mathfrak{g}$ in V , and denote by Π_λ the projection in V onto the isotypic component of V_λ parallel to $\bigoplus_{\substack{\mu \in P_+^\mathbb{L} \\ \mu \neq \lambda}} V_\mu$.

It suffices to prove existence and uniqueness of an extension $\tilde{\pi}$ of π to $F(\mathfrak{g}, \mathfrak{l})_q$ that possesses the following properties:

- i) $\tilde{\pi}(\mathcal{P}_\lambda) = \Pi_\lambda$, $\lambda \in P_+^\mathbb{L}$,
- ii) $\tilde{\pi}|_{R(\mathfrak{g}, \mathfrak{l})_q}$ is an approximately unital representation of $R(\mathfrak{g}, \mathfrak{l})_q$,
- iii) $\pi(\xi)v = \lim_{\Lambda \uparrow P_+^\mathbb{L}} \tilde{\pi}(\xi \chi_\Lambda)v$, $v \in V$, $\xi \in U_q \mathfrak{g}$.

Uniqueness of such extension is obvious. To prove its existence, we consider subsequently the following cases:

1. $V = V^{\text{univ}}$;
2. $V = \tilde{V}^{\text{univ}}$, with $\tilde{V}^{\text{univ}}$ being a multiple of the $U_q\mathfrak{g}$ -module V^{univ} ;
3. V is a submodule of $\tilde{V}^{\text{univ}}$;
4. V is a standard projective object in $C(\mathfrak{g}, \mathfrak{l})_q$;
5. (the general case) $V \in C(\mathfrak{g}, \mathfrak{l})_q$.

In the case 1) the desired statement follows from the definition of $R(\mathfrak{g}, \mathfrak{l})_q$. A passage from 1) to 2) and to 3) is obvious. To pass from 3) to 4), it is sufficient to use the fact that every $U_q\mathfrak{g}$ -module $P(V)$, $V \in C(\mathfrak{l}, \mathfrak{l})_q$, admits an embedding into a multiple $\tilde{V}^{\text{univ}}$ of the $U_q\mathfrak{g}$ -module V^{univ} . This is because every locally finite dimensional $U_q\mathfrak{l}$ -module admits an embedding into a $U_q\mathfrak{l}$ -module $\tilde{L}^{\text{univ}}$, which is a multiple of L^{univ} and has the property $P(\tilde{L}^{\text{univ}}) = \tilde{V}^{\text{univ}}$. Finally, to pass from 4) to 5), one can apply the existence of a surjective morphism $P(V) \rightarrow V$ in the category $C(\mathfrak{g}, \mathfrak{l})_q$. $\square$

Recall that the Weyl group W is generated by simple reflections $s_1, s_2, \dots, s_l$ and acts on the weight lattice P .

Proposition 2.7 1. *There exists a unique one-dimensional representation $\tilde{\varepsilon}$ of $F(\mathfrak{g}, \mathfrak{l})_q$ with*

$$\tilde{\varepsilon}(\xi) = \varepsilon(\xi), \quad \xi \in U_q\mathfrak{g}; \quad \tilde{\varepsilon}(P_\lambda) = \begin{cases} 1, & \lambda = 0, \\ 0, & \lambda \in P_+^{\mathbb{L}} \setminus \{0\}. \end{cases}$$

2. *There exists a unique anti-automorphism $\tilde{S}$ of the algebra $F(\mathfrak{g}, \mathfrak{l})_q$ such that*

$$\tilde{S}(\xi) = S(\xi), \quad \xi \in U_q\mathfrak{g}; \quad \tilde{S}(P_\lambda) = P_{-w_0^{\mathbb{L}}\lambda}, \quad \lambda \in P_+^{\mathbb{L}},$$

where $w_0^{\mathbb{L}}$ is the longest element of the Weyl group $W_{\mathbb{L}} \subset W$ generated by the simple reflections s_i , $i \in \mathbb{L}$.

Proof. Uniqueness of $\tilde{\varepsilon}$ and $\tilde{S}$ is obvious. Existence of $\tilde{\varepsilon}$ has been demonstrated in the proof of Theorem 2.6. Turn to proving existence of $\tilde{S}$. In the special case $\mathbb{G} = \mathbb{L}$ it follows from the definition of the algebra $F(\mathfrak{l})_q \hookrightarrow \text{End } L^{\text{univ}}$ and the well-known isomorphism [2, p. 168]

$$L(\mathfrak{l}, \lambda)^* \xrightarrow{\sim} L(\mathfrak{l}, -w_0^{\mathbb{L}}\lambda), \quad v(\mathfrak{l}, \lambda) \mapsto v(\mathfrak{l}, w_0^{\mathbb{L}}\lambda).$$

To pass from the special case $\mathbb{G} = \mathbb{L}$ to the general case $\mathbb{G} \supset \mathbb{L}$, consider the algebra $F(\mathfrak{g}, \mathfrak{l})_q^{\text{op}}$ which is derived from $F(\mathfrak{g}, \mathfrak{l})_q$ by replacement of the multiplication by the opposite one. $U_q\mathfrak{g}$ can be embedded into $F(\mathfrak{g}, \mathfrak{l})_q^{\text{op}}$:

$$U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q^{\text{op}}, \quad \xi \mapsto S(\xi).$$

The algebra $F(\mathfrak{l})_q$ also admits an embedding into $F(\mathfrak{g}, \mathfrak{l})_q^{\text{op}}$:

$$U_q\mathfrak{l} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q^{\text{op}}, \quad \xi \mapsto S(\xi); \quad P_\lambda \mapsto P_{-w_0^{\mathbb{L}}\lambda}.$$

It is easy to verify (F1) – (F4) for this pair of embeddings. What remains is to use the uniqueness of such pair, Proposition 2.4. $\square$

The category $C(\mathfrak{g}, \mathfrak{l})_q$ is canonically isomorphic to the category of approximately unital $R(\mathfrak{g}, \mathfrak{l})_q$ -modules. In what follows we identify these categories.

Consider the Hopf subalgebra $U_q \mathfrak{q}_{\mathbb{L}}^+ \subset U_q \mathfrak{g}$ generated by

$$E_i, \quad i \in \mathbb{G}; \quad F_j, \quad j \in \mathbb{L}; \quad K_m^{\pm 1}, \quad m = 1, 2, \dots, l.$$

and the Hopf subalgebra $U_q \mathfrak{q}_{\mathbb{L}}^- \subset U_q \mathfrak{g}$ generated by

$$E_i, \quad i \in \mathbb{L}; \quad F_j, \quad j \in \mathbb{G}; \quad K_m^{\pm 1}, \quad m = 1, 2, \dots, l.$$

The following statement is well known in the classical case $q = 1$ (see [7, p. 90]).

Proposition 2.8 *The linear maps*

$$U_q \mathfrak{g} \otimes_{U_q \mathfrak{l}} R(\mathfrak{l})_q \rightarrow R(\mathfrak{g}, \mathfrak{l})_q, \quad \xi \otimes r \mapsto \xi r, \quad (2.2)$$

$$R(\mathfrak{l})_q \otimes_{U_q \mathfrak{l}} U_q \mathfrak{g} \rightarrow R(\mathfrak{g}, \mathfrak{l})_q, \quad r \otimes \xi \mapsto r \xi,$$

are bijective.

Proof. By Proposition 2.7, it suffices to consider the linear map (2.2). Let us prove that it is onto. Note that for any $\lambda \in P_+^{\mathbb{L}}$ and any finite subset $\Lambda \subset P_+^{\mathbb{L}}$ containing λ there exists an element z in the center $Z(U_q \mathfrak{l})$ of the algebra $U_q \mathfrak{l}$ such that $P_\lambda = z \chi_\Lambda$. On the other hand, for any $\lambda \in P_+^{\mathbb{L}}$, $\xi \in U_q \mathfrak{g}$ there exist finite subsets $\Lambda', \Lambda'' \ni \lambda$ such that

$$P_\lambda \xi = P_\lambda \xi \chi_{\Lambda'}, \quad \xi \chi_{\Lambda'} = \chi_{\Lambda''} \xi \chi_{\Lambda'}.$$

Hence $P_\lambda \xi = (P_\lambda \chi_{\Lambda''}) \xi \chi_{\Lambda'} = z \chi_{\Lambda''} \xi \chi_{\Lambda'} = z \xi \chi_{\Lambda'}$. Thus for any $\xi \in U_q \mathfrak{g}$ and any finite subset $\Lambda \subset P_+^{\mathbb{L}}$ there exist such $\tilde{\xi} \in U_q \mathfrak{g}$ and a finite subset $\tilde{\Lambda} \subset P_+^{\mathbb{L}}$ that $\chi_\Lambda \xi = \tilde{\xi} \chi_{\tilde{\Lambda}}$. This can be readily used to prove that the linear map (2.2) is onto.

Prove that it is injective. One can use well-known results on bases in quantum universal enveloping algebras [6, Chapter 8] to derive a decomposition $U_q \mathfrak{g} = U_q \mathfrak{q}_{\mathbb{L}}^- \otimes_{U_q \mathfrak{l}} U_q \mathfrak{q}_{\mathbb{L}}^+$. It follows from this decomposition and Lemma A.1 that $U_q \mathfrak{g}$ is a free right $U_q \mathfrak{l}$ -module. Choose a free basis $\{e_i\}$. Let a be a non-zero element of $U_q \mathfrak{g} \otimes_{U_q \mathfrak{l}} R(\mathfrak{l})_q$. It has the form

$$a = \sum_i e_i \otimes r_i, \quad r_i \in R(\mathfrak{l})_q.$$

Let $\tilde{a}$ be the image of a under the map (2.2). We treat it as an element of $\text{End } V^{\text{univ}}$. It suffices to prove that the restriction of the linear map $\tilde{a}$ to the subspace $\{1 \otimes v \mid v \in L^{\text{univ}}\}$ is non-zero. It is easy to verify that $\tilde{a}(1 \otimes v) = \sum_i e_i \otimes r_i v$ for any $v \in L^{\text{univ}}$. It remains to use our choice of $\{e_i\}$ and the fact that the representation of $U_q \mathfrak{l}$ in L^{univ} is faithful. $\square$

3 The functors ind and Π

Consider two pairs of subsets

$$\mathbb{L} \subset \mathbb{G} \subset \{1, 2, \dots, l\}, \quad \mathbb{L}_1 \subset \mathbb{G}_1 \subset \{1, 2, \dots, l\},$$

with $\mathbb{L}_1 \subset \mathbb{L}$, $\mathbb{G}_1 \subset \mathbb{G}$. Obviously, one has embeddings of the associated Hopf subalgebras

$$U_q \mathfrak{l}_1 \subset U_q \mathfrak{l}, \quad U_q \mathfrak{g}_1 \subset U_q \mathfrak{g}.$$

$R(\mathfrak{g}, \mathfrak{l})_q$ is a left ideal in $F(\mathfrak{g}, \mathfrak{l})_q$ and hence is a left $U_q \mathfrak{g}_1$ -module. We claim that $R(\mathfrak{g}, \mathfrak{l})_q$ is an approximately unital left $R(\mathfrak{g}_1, \mathfrak{l}_1)_q$ -module. In fact, $R(\mathfrak{g}, \mathfrak{l})_q$ is an approximately unital left $R(\mathfrak{g}, \mathfrak{l})_q$ -module, hence a $U_q \mathfrak{g}$ -module of the category $C(\mathfrak{g}, \mathfrak{l})_q$ given by (1.1). Since $U_q \mathfrak{l}_1 \subset U_q \mathfrak{l}$, the module $R(\mathfrak{g}, \mathfrak{l})_q$ is also in the category $C(\mathfrak{g}, \mathfrak{l}_1)_q$. Finally, in view of $U_q \mathfrak{g}_1 \subset U_q \mathfrak{g}$, we are inside the category $C(\mathfrak{g}_1, \mathfrak{l}_1)_q$, which is equivalent to our claim. In a similar way, one can prove that $R(\mathfrak{g}, \mathfrak{l})_q$ is a right $R(\mathfrak{g}_1, \mathfrak{l}_1)_q$ -module.⁶

Throughout the rest of this section we follow [7] and replace the groups involved therein with the corresponding quantum universal enveloping algebras. Introduce the functor $P_{\mathfrak{g}_1, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}$ from the category $C(\mathfrak{g}_1, \mathfrak{l}_1)_q$ to the category $C(\mathfrak{g}, \mathfrak{l})_q$ by defining it on objects from $C(\mathfrak{g}_1, \mathfrak{l}_1)_q$ as follows:

$$P_{\mathfrak{g}_1, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}(Z) = R(\mathfrak{g}, \mathfrak{l})_q \otimes_{R(\mathfrak{g}_1, \mathfrak{l}_1)_q} Z.$$

The action on morphisms is defined in an obvious way.

$C(\mathfrak{g}_1, \mathfrak{l}_1)_q$ has enough projectives and the functor $P_{\mathfrak{g}_1, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}$ is covariant and right exact.⁷ Hence one has well defined derived functors: $\left(P_{\mathfrak{g}_1, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}\right)_j$, $j \in \mathbb{Z}_+$, from $C(\mathfrak{g}_1, \mathfrak{l}_1)_q$ to $C(\mathfrak{g}, \mathfrak{l})_q$.

Consider the two special cases: $\mathfrak{l}_1 = \mathfrak{l}$ and $\mathfrak{g}_1 = \mathfrak{g}$. Start from the first one. Let $\text{ind}_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}$ be the functor from the category $C(\mathfrak{g}_1, \mathfrak{l})_q$ to the category $C(\mathfrak{g}, \mathfrak{l})_q$ defined on objects as follows:

$$\text{ind}_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}(Z) = U_q \mathfrak{g} \otimes_{U_q \mathfrak{g}_1} Z.$$

The action on morphisms is defined in an obvious way. Just as in the classical case $q = 1$, one gets an isomorphism of functors $P_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}$ and $\text{ind}_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}$. Describe briefly this construction. The functor $\text{ind}_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}$ is left adjoint to the forgetful functor $\mathcal{F}_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}} : C(\mathfrak{g}, \mathfrak{l})_q \rightarrow C(\mathfrak{g}_1, \mathfrak{l})_q$ determined by the embedding $U_q \mathfrak{g}_1 \rightarrow U_q \mathfrak{g}$. On the other hand, $P_{\mathfrak{g}_1, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}$ is left adjoint to the functor $(\mathcal{F}^\vee)_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}}$, to be defined below (cf. [7, Proposition 2.34]). The functor $(\mathcal{F}^\vee)_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}} : C(\mathfrak{g}, \mathfrak{l})_q \rightarrow C(\mathfrak{g}_1, \mathfrak{l})_q$ is defined on objects of the category $C(\mathfrak{g}, \mathfrak{l})_q$ as follows:

$$(\mathcal{F}^\vee)_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}}(X) = \text{Hom}_{R(\mathfrak{g}, \mathfrak{l})_q}(R(\mathfrak{g}, \mathfrak{l})_q, X)_{\mathfrak{l}}.$$

A structure of $U_q \mathfrak{g}_1$ -module in $\text{Hom}_{R(\mathfrak{g}, \mathfrak{l})_q}(R(\mathfrak{g}, \mathfrak{l})_q, X)$ is imposed via the structure of right $U_q \mathfrak{g}_1$ -module in $R(\mathfrak{g}, \mathfrak{l})_q$, and the subscript $\mathfrak{l}$ stands for distinguishing the maximal submodule in the category $C(\mathfrak{g}_1, \mathfrak{l})_q$. The action of $(\mathcal{F}^\vee)_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}}$ on morphisms is defined in an obvious way. What remains is to construct an isomorphism of functors $\mathcal{F}_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}} \xrightarrow{\sim} (\mathcal{F}^\vee)_{\mathfrak{g}, \mathfrak{l}}^{\mathfrak{g}_1, \mathfrak{l}}$ (cf. [7, Proposition 2.33]). Let $X \in C(\mathfrak{g}, \mathfrak{l})_q$ and $x \in X$. Associate to every x a morphism of $R(\mathfrak{g}, \mathfrak{l})_q$ -modules given by

$$R(\mathfrak{g}, \mathfrak{l})_q \rightarrow X, \quad r \mapsto rx.$$

One can verify that the map $X \rightarrow \text{Hom}_{R(\mathfrak{g}, \mathfrak{l})_q}(R(\mathfrak{g}, \mathfrak{l})_q, X)$, which arises this way, is a morphism of $U_q \mathfrak{g}_1$ -modules and provides the desired isomorphism of functors.

An important consequence is the observation that for any $V \in C(\mathfrak{l}, \mathfrak{l})_q$ the standard projective object $P(V)$ in the category $C(\mathfrak{g}, \mathfrak{l})_q$ is canonically isomorphic to $P_{\mathfrak{l}, \mathfrak{l}}^{\mathfrak{g}, \mathfrak{l}}(V)$.

In the second special case $\mathfrak{g}_1 = \mathfrak{g}$ the functor in question is called the Bernstein functor and it is denoted by $\Pi_{\mathfrak{g}, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}$:

$$\Pi \equiv \Pi_{\mathfrak{g}, \mathfrak{l}_1}^{\mathfrak{g}, \mathfrak{l}}(Z) = R(\mathfrak{g}, \mathfrak{l})_q \otimes_{R(\mathfrak{g}, \mathfrak{l}_1)_q} Z, \quad Z \in C(\mathfrak{g}_1, \mathfrak{l}_1)_q.$$

⁶One can see from the proof that $R(\mathfrak{g}, \mathfrak{l})_q$ is a $R(\mathfrak{g}_1, \mathfrak{l}_1)_q$ -bimodule.

⁷See [7, p. 840]

In the classical case $q = 1$ the derived functors Π_j are crucial in constructing unitarizable Harish-Chandra modules via the Vogan-Zuckerman cohomological induction [7]. Turn to describing a quantum analogue for this method.

4 A quantum analogue for cohomological induction

We assume in the sequel that $\mathbb{G} = \{1, 2, \dots, l\}$, and hence $U_q \mathfrak{g} = U_q$.

Recall (see [4, 8]) a definition of the pairs $(\mathfrak{g}, \mathfrak{k})$ used in a construction of bounded symmetric domains via the Harish-Chandra embedding. Here $\mathfrak{g}$ is a simple complex Lie algebra of type A, B, C, D, E_6, E_7 . A Hopf subalgebra $U_q \mathfrak{k} \subset U_q \mathfrak{g}$ is generated by $E_i, F_i, i \in \mathbb{K}; K_j^{\pm 1}, j = 1, 2, \dots, l$, with $\mathbb{K} = \{1, 2, \dots, l\} \setminus \{l_0\}$. We assume also that the simple root α_{l_0} has coefficient 1 in the decomposition of the maximal root

$$\delta = \sum_{i=1}^l n_i \alpha_i, \quad n_{l_0} = 1. \quad (4.1)$$

In what follows the notation $(\mathfrak{g}, \mathfrak{k})$ will stand only for such pairs.

To clarify the precise nature of condition (4.1), equip the Lie algebra $\mathfrak{g}$ with a grading as follows: $\deg E_{l_0} = 1, \deg F_{l_0} = -1, \deg H_{l_0} = 0, \deg E_j = \deg F_j = \deg H_j = 0$ for $j \neq l_0$. The automorphism θ of $\mathfrak{g}$ given by the formula

$$\theta(\xi) = (-1)^{\deg \xi} \xi$$

is an involution. Now (4.1) implies that

$$\mathfrak{g} = \mathfrak{p}^- \oplus \mathfrak{k} \oplus \mathfrak{p}^+, \quad \mathfrak{p}^\pm = \{\xi \in \mathfrak{g} \mid \deg \xi = \pm 1\}.$$

It is worthwhile to note that the pairs $(\mathfrak{g}, \mathfrak{k})$ in question are complexifications of the pairs $(\mathfrak{g}_0, \mathfrak{k}_0)$, with $\mathfrak{g}_0$ being the Lie subalgebra of the automorphism group of an irreducible bounded symmetric domain, and $\mathfrak{k}_0$ being the Lie algebra of the stabilizer of a point in this domain [4].

Equip $U_q \mathfrak{g}$ with a structure of Hopf $*$ -algebra via the involution $*$ defined as follows:

$$E_j^* = \begin{cases} -K_j F_j, & j = l_0; \\ K_j F_j, & j \neq l_0; \end{cases} \quad F_j^* = \begin{cases} -E_j K_j^{-1}, & j = l_0; \\ E_j K_j^{-1}, & j \neq l_0; \end{cases}$$

$$(K_j^{\pm 1})^* = K_j^{\pm 1}, \quad j = 1, 2, \dots, l.$$

A module V over a Hopf $*$ -algebra A is said to be unitarizable if it admits a positive definite invariant form $(\cdot, \cdot)$:

$$(av_1, v_2) = (v_1, a^* v_2), \quad a \in A, v_1, v_2 \in V.$$

The cohomological induction is among the tools for constructing unitarizable modules of the category $C(\mathfrak{g}, \mathfrak{k})$ in the case $q = 1$. Describe a q -analogue for this method.

Let $\mathbb{L} \subset \{1, 2, \dots, l\}$, $\mathbb{L} \not\subset \mathbb{K}$, and $U_q \mathfrak{l}$ is the Hopf subalgebra corresponding to the subset $\mathbb{L}$. Obviously, $U_q \mathfrak{l}$ inherits the structure of Hopf $*$ -algebra. We intend to derive a unitarizable $U_q \mathfrak{g}$ -module of the category $C(\mathfrak{g}, \mathfrak{k})_q$ starting from a unitarizable $U_q \mathfrak{l}$ -module of the category $C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$.⁸

⁸ $U_q(\mathfrak{l} \cap \mathfrak{k})$ is a Hopf subalgebra corresponding to the subset $\mathbb{L} \cap \mathbb{K}$.

If $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P$ and $\lambda_j = 0$ for all $j \in \mathbb{L}$, one has the following well defined one-dimensional $U_q \mathfrak{l}$ -module $\mathbb{C}_\lambda$:

$$E_j \mathbf{1} = F_j \mathbf{1} = 0, \quad j \in \mathbb{L}; \quad K_j^{\pm 1} \mathbf{1} = q_j^{\pm \lambda_j} \mathbf{1}, \quad j = 1, 2, \dots, l.$$

As an important example of such linear functional one should mention the difference $\rho_u = \rho - \rho_l$ between the half-sum ρ of positive roots of the Lie algebra $\mathfrak{g}$ and the half-sum ρ_l of positive roots of the Lie algebra $\mathfrak{l}$:

$$\rho_u(H_j) = \begin{cases} 1, & j \notin \mathbb{L}, \\ 0, & j \in \mathbb{L}. \end{cases}$$

Every $U_q \mathfrak{l}$ -module $Z \in C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$ determines a $U_q \mathfrak{l}$ -module $Z^\# = Z \otimes \mathbb{C}_{2\rho_u} \in C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$. Equip it with a structure of $U_q \mathfrak{q}_\mathbb{L}^-$ -module via the surjective morphism of algebras $U_q \mathfrak{q}_\mathbb{L}^- \rightarrow U_q \mathfrak{l}$:

$$F_i \mapsto \begin{cases} F_i, & i \in \mathbb{L} \\ 0, & i \in \{1, 2, \dots, l\} \setminus \mathbb{L} \end{cases}, \quad E_j \mapsto E_j, \quad K_j^{\pm 1} \mapsto K_j^{\pm 1}, \quad j \in \{1, 2, \dots, l\}.$$

The generalized Verma module $\text{ind}_{\mathfrak{q}_\mathbb{L}^-}^{\mathfrak{g}} Z^\# \stackrel{\text{def}}{=} U_q \mathfrak{g} \otimes_{U_q \mathfrak{q}_\mathbb{L}^-} Z^\#$ belongs to the category $C(\mathfrak{g}, \mathfrak{l})_q$ (see Lemma A.1). Hence the equality

$$\mathcal{L}_j(Z) = \left(\Pi_{\mathfrak{g}, \mathfrak{l} \cap \mathfrak{k}}^{\mathfrak{g}, \mathfrak{k}} \right)_j \left(\text{ind}_{\mathfrak{q}_\mathbb{L}^-}^{\mathfrak{g}} (Z^\#) \right)$$

determines functors $\mathcal{L}_j$, $j \in \mathbb{Z}_+$, from $C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$ to $C(\mathfrak{g}, \mathfrak{k})_q$.

In the classical case $q = 1$ under suitable dominance assumptions on Z the only non-zero modules $\mathcal{L}_s(Z)$ are those with $s = \frac{1}{2}(\dim \mathfrak{k} - \dim(\mathfrak{k} \cap \mathfrak{l}))$ [7, p. 369]. So, a particular interest is in considering the $U_q \mathfrak{g}$ -modules

$$A_q(\lambda) \stackrel{\text{def}}{=} \mathcal{L}_s(\mathbb{C}_\lambda), \quad s = \frac{1}{2}(\dim \mathfrak{k} - \dim(\mathfrak{k} \cap \mathfrak{l})),$$

with $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P$ and $\lambda_j = 0$ for $j \in \mathbb{L}$.

Appendix. Proofs of the Lemmas

Let $\mathbb{G} \supset \mathbb{L}$ be subsets of $\{1, 2, \dots, l\}$. Equip $U_q \mathfrak{g}$ with a grading by setting

$$\deg E_j = \begin{cases} 1, & j \in \mathbb{G} \setminus \mathbb{L} \\ 0, & j \in \mathbb{L} \end{cases}, \quad \deg F_j = \begin{cases} -1, & j \in \mathbb{G} \setminus \mathbb{L} \\ 0, & j \in \mathbb{L} \end{cases},$$

$$\deg K_j^{\pm 1} = 0, \quad j = 1, 2, \dots, l.$$

Let $(U_q \mathfrak{q}_\mathbb{L}^\pm)_j = \{ \xi \in U_q \mathfrak{q}_\mathbb{L}^\pm \mid \deg \xi = j \}$. Obviously, $(U_q \mathfrak{q}_\mathbb{L}^\pm)_0 = U_q \mathfrak{l}$.

Lemma A.1 *The homogeneous components $(U_q \mathfrak{q}_\mathbb{L}^\pm)_j$ are free left and free right $U_q \mathfrak{l}$ -modules of finite rank.*

Proof. Recall that the subsets $\mathbb{L} \subset \mathbb{G} \subset \{1, 2, \dots, l\}$ determine the subgroups $W_{\mathbb{L}} \subset W_{\mathbb{G}}$ of the Weyl group W . Those subgroups are exactly the Weyl groups of the Lie algebras $\mathfrak{l}$ and $\mathfrak{g}$, respectively. Let $w_{0,\mathbb{L}}$ (respectively, $w_{0,\mathbb{G}}$) be the longest element of the Weyl group $W_{\mathbb{L}}$ (respectively, $W_{\mathbb{G}}$).

We start from the larger subset $\mathbb{G}$. Choose a reduced expression

$$w_{0,\mathbb{G}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_M}, \quad (\text{A.2})$$

with s_{i_k} being the reflection corresponding to the simple root α_{i_k} . Consider the Lusztig automorphisms T_i , $i \in \mathbb{G}$, of the algebra $U_q \mathfrak{g}'$ generated by E_i , F_i , $K_i^{\pm 1}$, $i \in \mathbb{G}$, [6, Chapter 8]. They have the form

$$\begin{aligned} T_i(K_j) &= K_j K_i^{-a_{ij}}, \\ T_i(E_i) &= -F_i K_i, \quad T_i(F_i) = -K_i^{-1} E_i, \\ T_i(E_j) &= \sum_{r+s=-a_{ij}} \text{const } E_i^s E_j E_i^r, \quad i \neq j, \\ T_i(F_j) &= \sum_{r+s=-a_{ij}} \text{const } F_i^r F_j F_i^s, \quad i \neq j. \end{aligned} \quad (\text{A.3})$$

In a similar way, consider the algebra $U_q \mathfrak{l}'$ associated to $\mathbb{L}$. Let $W^{\mathbb{L}} = \{w \in W_{\mathbb{G}} \mid l(ws) > l(w) \text{ for all simple reflections } s \in W_{\mathbb{L}}\}$, where $l(w)$ stands for the length of w . One has $w_{0,\mathbb{G}} = w^{\mathbb{L}} w_{0,\mathbb{L}}$ with $w^{\mathbb{L}} \in W^{\mathbb{L}}$ and $l(w_{0,\mathbb{G}}) = l(w^{\mathbb{L}}) + l(w_{0,\mathbb{L}})$ [5, Proposition 1.10(c)]. This allows one to rewrite the reduced expression (A.2) in the form

$$w_{0,\mathbb{G}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_{M'}} s_{i_{M'+1}} s_{i_{M'+2}} \dots s_{i_M}, \quad (\text{A.4})$$

with $w_{0,\mathbb{L}} = s_{i_{M'+1}} s_{i_{M'+2}} \dots s_{i_M}$ and $w^{\mathbb{L}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_{M'}}$.

We are about to apply the Lusztig theorem [6, Theorem 8.24]. Observe that the special reduced expression (A.4), in view of the explicit form of the Lusztig automorphisms (A.3) implies that all the monomials

$$T_{i_1} T_{i_2} \dots T_{i_{M'-1}}(E_{i_{M'}}^{a_{M'}}) \cdot T_{i_1} T_{i_2} \dots T_{i_{M'-2}}(E_{i_{M'-1}}^{a_{M'-1}}) \cdot \dots \cdot T_{i_1} T_{i_2}(E_{i_3}^{a_3}) \cdot T_{i_1}(E_{i_2}^{a_2}) \cdot E_{i_1}^{a_1}$$

are in $U_q \mathfrak{l}'$. It follows from the Lusztig theorem [6, Theorem 8.24] that the monomials

$$\begin{aligned} T_{i_1} T_{i_2} \dots T_{i_{M-1}}(E_{i_M}^{a_M}) \cdot T_{i_1} T_{i_2} \dots T_{i_{M-2}}(E_{i_{M-1}}^{a_{M-1}}) \cdot \dots \\ \dots \cdot T_{i_1} T_{i_2} \dots T_{i_{M'+1}}(E_{i_{M'+2}}^{a_{M'+2}}) \cdot T_{i_1} T_{i_2} \dots T_{i_{M'}}(E_{i_{M'+1}}^{a_{M'+1}}) \end{aligned} \quad (\text{A.5})$$

with all $a_{i_k} \in \mathbb{Z}_+$, form a free basis in the right $U_q \mathfrak{l}$ -module $U_q \mathfrak{q}_{\mathbb{L}}^+$. As one can readily separate out for each $j \in \mathbb{Z}_+$ finitely many such monomials that span the j -th homogeneous component, we thus get our claim for $(U_q \mathfrak{q}_{\mathbb{L}}^+)_j$ as a right $U_q \mathfrak{l}$ -module. All other claims can be proved in a similar way. $\square$

Corollary A.2 *Every $U_q \mathfrak{g}$ -module V contains the largest submodule $V_{\mathfrak{l}}$ of the category $C(\mathfrak{g}, \mathfrak{l})_q$.*

Proof. Obviously, V possesses the largest weight submodule $V_{\mathfrak{h}}$. What remains is to prove that the subspace $V_{\mathfrak{l}} = \{v \in V_{\mathfrak{h}} \mid \dim(U_q \mathfrak{l}v) < \infty\}$ is a submodule of the $U_q \mathfrak{g}$ -module V . Let $\xi \in (U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j$, $v \in V_{\mathfrak{l}}$. It follows from Lemma A.1 that for some $\{\eta_1, \eta_2, \dots, \eta_{N(j)}\}$

$$(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j = \sum_{k=1}^{N(j)} \eta_k U_q \mathfrak{l}.$$

Hence,

$$\dim(U_q \mathfrak{l} \xi v) \leq \dim \left((U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j v \right) = \dim \left(\sum_{k=1}^{N(j)} \eta_k U_q \mathfrak{l} v \right) < \infty.$$

Thus $\xi v \in V_{\mathfrak{l}}$. □

Corollary A.3 *For any $V \in C(\mathfrak{l}, \mathfrak{l})_q$, the module $P(V)$ belongs to the category $C(\mathfrak{g}, \mathfrak{l})_q$.*

Proof. Since $U_q \mathfrak{g}$ is a free right $U_q \mathfrak{l}$ -module, one has an embedding of $U_q \mathfrak{l}$ -modules

$$i : V \hookrightarrow P(V), \quad i : v \mapsto 1 \otimes v.$$

The relation $P(V) = P(V)_{\mathfrak{l}}$ is due to

$$P(V) = U_q \mathfrak{g} i(V) \subset U_q \mathfrak{g} P(V)_{\mathfrak{l}} = P(V)_{\mathfrak{l}}. \quad \square$$

The rest of the statements of Lemma 2.1 can be proved in the same way as in the classical case $q = 1$.

To prove Lemma 2.5 we need the following auxiliary statement.

Lemma A.4 *Let $\lambda \in P_+^{\mathbb{L}}$ and $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j$ be the homogeneous components of the graded algebras $U_q \mathfrak{q}_{\mathbb{L}}^{\pm}$.*

1. *The vector spaces $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathcal{P}_{\lambda} \subset \text{End } V^{\text{univ}}$ are finite dimensional.*
2. *The vector spaces $\mathcal{P}_{\lambda} \cdot (U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j \subset \text{End } V^{\text{univ}}$ are finite dimensional.⁹*

Proof. Prove the first statement. We consider $\text{End } V^{\text{univ}}$ as a $U_q \mathfrak{l}$ -module with respect to the action as follows:

$$(\xi a) : v \mapsto \xi(av), \quad v \in V^{\text{univ}}, \quad a \in \text{End } V^{\text{univ}}, \quad \xi \in U_q \mathfrak{l}.$$

Obviously, $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathcal{P}_{\lambda}$ is a submodule of the $U_q \mathfrak{l}$ -module $\text{End } V^{\text{univ}}$.

Let $\pi_{\lambda} : U_q \mathfrak{l} \rightarrow \text{End } L(\mathfrak{l}, \lambda)$ be the representation of the Hopf subalgebra $U_q \mathfrak{l}$ corresponding to the $U_q \mathfrak{l}$ -module $L(\mathfrak{l}, \lambda)$. If $\xi \in \text{Ker } \pi_{\lambda}$, then $\xi \cdot \mathcal{P}_{\lambda} = 0$. Hence the diagram

$$\begin{array}{ccc} U_q \mathfrak{l} & \xrightarrow{\xi \mapsto \xi \cdot \mathcal{P}_{\lambda}} & \text{End } V^{\text{univ}} \\ \pi_{\lambda} \downarrow & \nearrow & \\ \text{End } L(\mathfrak{l}, \lambda) & & \end{array}$$

⁹A dot is used here to denote the product of elements in $\text{End } V^{\text{univ}}$.

can be completed up to a commutative one, and

$$\dim(U_q \mathfrak{l} \cdot \mathcal{P}_\lambda) \leq (\dim(L(\mathfrak{l}, \lambda)))^2.$$

Thus the first statement of the Lemma is proved in the case $j = 0$. It remains to elaborate the fact that $(U_q \mathfrak{q}_\mathbb{L}^\pm)_j$ is a right $U_q \mathfrak{l}$ -module of finite rank (see Lemma A.1).

Now the first statement is proved. The second one can be proved in a similar way using the commutative diagram

$$\begin{array}{ccc} U_q \mathfrak{l} & \xrightarrow{\xi \mapsto \mathcal{P}_\lambda \cdot \xi} & \text{End } V^{\text{univ}} \\ \pi_\lambda \downarrow & \nearrow & \\ \text{End } L(\mathfrak{l}, \lambda) & & \end{array}$$

in the category of $U_q \mathfrak{l}^{\text{op}}$ -modules.¹⁰ □

Turn to the proof of Lemma 2.5. Consider the subspace $A \subset U_q \mathfrak{g}$ of such elements $\xi \in U_q \mathfrak{g}$ that for any $\lambda \in P_+^\mathbb{L}$ there exists a finite subset $\Lambda \subset P_+^\mathbb{L}$ with

$$\xi \cdot \mathcal{P}_\lambda = \chi_\Lambda \cdot \xi \cdot \mathcal{P}_\lambda, \quad (\text{A.6})$$

$$\mathcal{P}_\lambda \cdot \xi = \mathcal{P}_\lambda \cdot \xi \cdot \chi_\Lambda. \quad (\text{A.7})$$

One has to prove that $A = U_q \mathfrak{g}$. For that, it suffices to demonstrate that A is a subalgebra of $U_q \mathfrak{g}$ and $A \supset (U_q \mathfrak{q}_\mathbb{L}^\pm)_j$ for all j .

Prove the first statement. It is obvious that A is a vector subspace. Now let $\xi, \eta \in A$, and consider the product $\xi\eta$. A double application of (A.6) first w.r.t. η and then ξ allows one to deduce that the image of the linear operator $\xi \cdot \eta \cdot \mathcal{P}_\lambda \in \text{End } V^{\text{univ}}$ is accommodated by the sum of finitely many isotypic components V_λ^{univ} . Let this sum be $\bigoplus_{\lambda' \in \Lambda'} V_{\lambda'}^{\text{univ}}$, then

$\xi \cdot \eta \cdot \mathcal{P}_\lambda = \chi_{\Lambda'} \cdot \xi \cdot \eta \cdot \mathcal{P}_\lambda$. Thus we get (A.6) for $\xi\eta$.

In a similar way, apply (A.7) twice to deduce that for some finite subset $\Lambda'' \subset P_+^\mathbb{L}$ one has $\text{Ker}(\mathcal{P}_\lambda \cdot \xi \cdot \eta) \supset (\text{id} - \chi_{\Lambda''}) V^{\text{univ}}$, hence $\mathcal{P}_\lambda \cdot \xi \cdot \eta = \mathcal{P}_\lambda \cdot \xi \cdot \eta \cdot \chi_{\Lambda''}$. Thus (A.7) holds for $\xi\eta$.

Turn to the second statement. We restrict ourselves to proving that (A.6) is valid for all $\xi \in (U_q \mathfrak{q}_\mathbb{L}^\pm)_j$. It follows from Lemma A.4 that for every j, λ , the vector space $(U_q \mathfrak{q}_\mathbb{L}^\pm)_j \cdot \mathcal{P}_\lambda$ is finite dimensional. It is also a $U_q \mathfrak{l}$ -module since $U_q \mathfrak{l} (U_q \mathfrak{q}_\mathbb{L}^\pm)_j \subset (U_q \mathfrak{q}_\mathbb{L}^\pm)_j$. Hence $(U_q \mathfrak{q}_\mathbb{L}^\pm)_j \cdot \mathcal{P}_\lambda$ is a sum of finitely many $U_q \mathfrak{l}$ -isotypic components. On the other hand, if for some $a \in \text{End } V^{\text{univ}}$, the $U_q \mathfrak{l}$ -module $U_q \mathfrak{l} \cdot a \subset \text{End } V^{\text{univ}}$ is a multiple of $L(\mathfrak{l}, \mu)$, $\mu \in P_+^\mathbb{L}$, one has $a V^{\text{univ}} \subset V_\mu^{\text{univ}}$ and $\mathcal{P}_\mu \cdot a = a$. It follows that $\mathcal{P}_{\lambda'} \cdot (U_q \mathfrak{q}_\mathbb{L}^\pm)_j \cdot \mathcal{P}_\lambda = 0$ for all but finitely many $\lambda' \in P_+^\mathbb{L}$. Hence $\xi \cdot \mathcal{P}_\lambda = \chi_\Lambda \cdot \xi \cdot \mathcal{P}_\lambda$ for some finite subset $\Lambda \subset P_+^\mathbb{L}$. □

References

- [1] R. J. Baston, M. G. Eastwood. The Penrose Transform. Its Interaction with Representation Theory. Clarendon Press, Oxford, 1989.
- [2] N. Bourbaki, Groupes et Algèbres de Lie, chapitres VII-VIII. Herman, 1975.

¹⁰The algebra $U_q \mathfrak{l}^{\text{op}}$ is derived from $U_q \mathfrak{l}$ by replacing the multiplication with the opposite one.

- [3] C. De Concini, V. Kac. *Representations of quantum groups at roots of 1*. In: Operator Algebras, Unitary Representations, Enveloping Algebras and Invariant Theory, A. Connes, M. Duflo, R. Rentschler (eds.), Birkhäuser, Boston-Basel-Berlin, 1990, 472 – 506.
- [4] S. Helgason. *Differential Geometry and Symmetric Spaces*. Academic Press, New York, London, 1962.
- [5] J. E. Humphreys. *Reflection Groups and Coxeter Groups*. Cambridge Univ. Press, Cambridge, 1990.
- [6] J. C. Jantzen. *Lectures on Quantum Groups*. American Mathematical Society, Providence, R. I., 1996.
- [7] A. Knapp, D. A. Vogan, Jr. *Cohomological Induction and Unitary Representations*. Princeton Univ. Press, Princeton, 1995.
- [8] A. Korányi, *Function spaces on bounded symmetric domains*, In: Analysis and geometry on complex homogeneous domains. Birkhäuser Boston, Inc., Boston, MA, 2000, J. Faraut, S. Kaneyuki, A. Koranyi, Qi-keng Lu, Guy Roos, (eds), 183 – 281.
- [9] D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman *Non-compact quantum groups and quantum Harish-Chandra modules*. In: Supersymmetry and Quantum Field Theory, North-Holland, D. Sorokin (ed.), the Netherlands, ser. Nucl. Phys. B (Proc. Suppl.), **102 & 103** (2001), 334 – 337.
- [10] D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman, *On a q -analogue of the Penrose transform*, Ukrainian Phys. J. **47** (2002), No 3, 288 – 292.
- [11] S. Sinel'shchikov, L. Vaksman, *On q -analogues of bounded symmetric domains and Dolbeault complexes*, Math. Phys. Anal. Geom. **1** (1998), No 1, 75 – 100.